



# Disabilities "Match-up"

High School

1. Social/communication disorder – difficulty with transitioning to “new things” – insistence on “sameness”
2. Disorder in psychological processing resulting in an impaired ability to read, write, do math, etc.
3. Significantly below average intellectual functioning coupled with delays in adaptive skills (i.e., skills of daily living).
4. Neurological disability causing hyperactivity and/or impulsivity which impairs the ability to pay attention in school.
5. A genetic disability impacting mostly boys, which is the 2<sup>nd</sup> most common inherited cause of mental retardation.
6. A neurological disorder which causes the person to have “tics,” which may include uncontrolled motoric movements.
7. A psychiatric disorder in which the person has an obsession with objects, patterns of objects, or motions.
8. A disability of the muscles and joints which causes the arms and/or legs to be floppy or tightly contracted.
9. A disability in which the person has impaired hearing or no functional hearing.
10. A disability in which the person has limited or no sight.
11. A genetic disability which results in short stature and unique facial and hand features, and mild to moderate retardation.
12. A psychiatric disorder that causes individuals to experience “cycling” of mood states, ranging from a highly alert, agitated state to, depression.
13. A psychiatric disorder in which the person has an altered perception of reality, which may include hearing voices or unreasonable thoughts.

14. A psychiatric disorder in which certain life settings cause unreasonable fears or feelings of panic.
15. A psychiatric disorder in which the person feels unreasonably “down” or “sad,” not caused by any actual events.
16. A disability in which the person may lose consciousness, fall, and experience tightening and contracting of the muscles.
17. A disability in which the person with normal intelligence has minor social difficulties with social interactions and unusual experiences.
18. A disability in which damage to the spinal cord causes limited use of legs or inability to walk.



# *Disabilities "Match-up" Answer Sheet*

- \_\_\_\_\_ Mental Retardation
- \_\_\_\_\_ Bipolar
- \_\_\_\_\_ Autism
- \_\_\_\_\_ Specific Learning Disability
- \_\_\_\_\_ Attention Deficit/Hyperactivity Disorder
- \_\_\_\_\_ Anxiety Disorder
- \_\_\_\_\_ Fragile X
- \_\_\_\_\_ Schizophrenia
- \_\_\_\_\_ Obsessive Compulsive Disorder
- \_\_\_\_\_ Cerebral Palsy
- \_\_\_\_\_ Visual Impairment
- \_\_\_\_\_ Tourette Syndrome
- \_\_\_\_\_ Deaf/Hard of Hearing
- \_\_\_\_\_ Depression
- \_\_\_\_\_ Asperger's Syndrome
- \_\_\_\_\_ Down Syndrome
- \_\_\_\_\_ Spina Bifida
- \_\_\_\_\_ Epilepsy/Seizure Disorder

**Answer Key Follows**



# *Disabilities "Match-up" Answer Key*

- 3 Mental Retardation
- 12 Bipolar
- 1 Autism
- 2 Specific Learning Disability
- 4 Attention Deficit/Hyperactivity Disorder
- 14 Anxiety Disorder
- 5 Fragile X
- 13 Schizophrenia
- 7 Obsessive Compulsive Disorder
- 8 Cerebral Palsy
- 10 Visual Impairment
- 6 Tourette Syndrome
- 9 Deaf/Hard of Hearing
- 15 Depression
- 17 Asperger's Syndrome
- 11 Down Syndrome
- 18 Spina Bifida
- 16 Epilepsy/Seizure Disorder